HQExoTM **Exosome-hUCMSC**

Catalog: Exo-SC05

PRODUCT INFORMATION

Name	HQExo™ Exosome-hUCMSC
Cat No.	Exo-SC05
Source	Exosome derived from Human Umbilical Cord Mesenchymal Stem Cell
Product Overview	Exosomes are small endosome derived lipid nanoparticles (30-160 nm in diameter) secreted by exocytosis by most living cells and contain specifical cargos, such as RNAs, lipids, and proteins. The cargos amount and con position of exosomes depend on the cell type from which they are released, which making them useful for bio marker discovery and functional characterization. Exosomes have been isolated from mesenchymal stem cells (MSCs), which have captured great attention in regenerative and translational medicine over a few decades due to their low immunogenicity, high biocompatibility, differentiation capacity, potent immunomodulatory proper ties, and their ability to be favorably cultured and manipulated. Exosomes transfer functional cargos like miRNA and mRNA molecules, peptides, proteins, cytokines and lipids from MSCs to the recipient cells, which contributes to the healing of necrotic tissues and organs or apoptotic cells HQExo TM exosomes derived from MSCs could use as positive controls for exosome isolation and functional research, such as ELISA, FACS, WB. With the unique capacity of proliferation and differentiation, exosomes derived from MSCs represented a great opportunity for regenerative therapeutic agents. For example, it has been reported that exosomes from cardiomyocy es or stem/progenitor cells can promote cardiac repair and regeneration. Lyophilization is useful for a long-ter m storage at 4°C, and frozen liquid should be kept at -20°C to -80°C. Ultracentrifugation and precipitation techniques are mainly used in exosome Isolation. It had been reported that both methods yielded extracellular vesic les in the size range of exosomes and included apoproteins, which can be used in downstream analyses. Nanop articles Tracking Analysis (NTA) is used for measuring exosome particles concentration, and WB or ELISA can be used in exosomal biomarkers analysis. Creative Biostructure standard exosome products guarantee higher purity and quality to meet our customer research.
Form	Frozen liquid
Concentration	1x10^10 particles/ml
Storage	Frozen liquid store at -20°C to -80°C. Recommended to avoid repeated freeze-and-thaw cycles.

45-1 Ramsey Road, Shirley, NY 11967, USA

Tel:1-631-317-1417 Fax:1-631-207-8356