



HQExoTM **Exosome-COLO1**

Catalog: Exo-CH16

PRODUCT INFORMATION

Exosome derived from human colon carcinoma (COLO1 cell line) Exosomes are nanosized vesicles (30-160 nm) secreted by exocytosis by most cell types and contain specargos, such as RNAs, lipids, and proteins. The cargos amount and composition of exosomes depend on I type from which they are released, which making them useful for biomarker discovery and functional cerization. Exosomes have been isolated from cancer cell lines (human and mouse), which helps understa or growth microenvironments. Exosome derived from enormous model human cancer cell lines to improstudies of tumor growth and invasion signaling pathways as well as how these tumor exosomes function tan insight into antitumor research. HQExo TM standard exosomes could use as positive controls for exosolation and functional research, such as ELISA, FACS, WB. Lyophilization is useful for a long-term stor 4°C, and frozen liquid should be kept at -20°C to -80°C. Ultracentrifugation and precipitation techniques ainly used in exosome Isolation. It had been reported that both methods yielded extracellular vesicles in the erange of exosomes and included apoproteins, which can be used in downstream analyses. Nanoparticle king Analysis (NTA) is used for measuring exosome particles concentration, and WB or ELISA can be the exosomal biomarkers analysis. Creative Biostructure standard exosome products guarantee higher purity utality to meet our customer research. Lyophilized powder/ frozen liquid. Reconstitute lyophilized exosome by adding deionized water for a deformal inal concentration. Centrifuge before opening to ensure exosomes are at bottom, resuspend exosomes by ing and/or vortex, please avoid bubbles. Centrifuge again and mix well for using.		
Exosome derived from human colon carcinoma (COLO1 cell line) Exosomes are nanosized vesicles (30-160 nm) secreted by exocytosis by most cell types and contain specargos, such as RNAs, lipids, and proteins. The cargos amount and composition of exosomes depend on 1 type from which they are released, which making them useful for biomarker discovery and functional cerization. Exosomes have been isolated from cancer cell lines (human and mouse), which helps understa or growth microenvironments. Exosome derived from enormous model human cancer cell lines to improstudies of tumor growth and invasion signaling pathways as well as how these tumor exosomes function tan insight into antitumor research. HQExo TM standard exosomes could use as positive controls for exosolation and functional research, such as ELISA, FACS, WB. Lyophilization is useful for a long-term store 4°C, and frozen liquid should be kept at -20°C to -80°C. Ultracentrifugation and precipitation techniques ainly used in exosome Isolation. It had been reported that both methods yielded extracellular vesicles in a crange of exosomes and included apoproteins, which can be used in downstream analyses. Nanoparticle king Analysis (NTA) is used for measuring exosome particles concentration, and WB or ELISA can be usexosomal biomarkers analysis. Creative Biostructure standard exosome products guarantee higher purity uality to meet our customer research. Lyophilized powder/ frozen liquid. Reconstitute lyophilized exosome by adding deionized water for a de inal concentration. Centrifuge before opening to ensure exosomes are at bottom, resuspend exosomes by ing and/or vortex, please avoid bubbles. Centrifuge again and mix well for using.	Name	HQExo™ Exosome-COLO1
Exosomes are nanosized vesicles (30-160 nm) secreted by exocytosis by most cell types and contain specargos, such as RNAs, lipids, and proteins. The cargos amount and composition of exosomes depend on I type from which they are released, which making them useful for biomarker discovery and functional cerization. Exosomes have been isolated from cancer cell lines (human and mouse), which helps understator growth microenvironments. Exosome derived from enormous model human cancer cell lines to improstudies of tumor growth and invasion signaling pathways as well as how these tumor exosomes function tan insight into antitumor research. HQExo TM standard exosomes could use as positive controls for exosolation and functional research, such as ELISA, FACS, WB. Lyophilization is useful for a long-term store. A considerable of the constitution of the constitutio	Cat No.	Exo-CH16
cargos, such as RNAs, lipids, and proteins. The cargos amount and composition of exosomes depend on I type from which they are released, which making them useful for biomarker discovery and functional cerization. Exosomes have been isolated from cancer cell lines (human and mouse), which helps understated or growth microenvironments. Exosome derived from enormous model human cancer cell lines to improse studies of tumor growth and invasion signaling pathways as well as how these tumor exosomes function to an insight into antitumor research. HQExo™ standard exosomes could use as positive controls for exost olation and functional research, such as ELISA, FACS, WB. Lyophilization is useful for a long-term store 4°C, and frozen liquid should be kept at -20°C to -80°C. Ultracentrifugation and precipitation techniques ainly used in exosome Isolation. It had been reported that both methods yielded extracellular vesicles in the erange of exosomes and included apoproteins, which can be used in downstream analyses. Nanoparticle king Analysis (NTA) is used for measuring exosome particles concentration, and WB or ELISA can be used exosomal biomarkers analysis. Creative Biostructure standard exosome products guarantee higher purity uality to meet our customer research. Lyophilized powder/ frozen liquid. Reconstitute lyophilized exosome by adding deionized water for a deformation in an analysis (NTA) is expected by the product of the	Source	Exosome derived from human colon carcinoma (COLO1 cell line)
inal concentration. Centrifuge before opening to ensure exosomes are at bottom, resuspend exosomes by ing and/or vortex, please avoid bubbles. Centrifuge again and mix well for using.	Product Overview	·
	Form	Lyophilized powder/ frozen liquid. Reconstitute lyophilized exosome by adding deionized water for a desired in in a concentration. Centrifuge before opening to ensure exosomes are at bottom, resuspend exosomes by pipetting and/or vortex, please avoid bubbles. Centrifuge again and mix well for using.
Concentration $> 1x10^8$ particles	Concentration	> 1x10^8 particles
Lyophilized powder store at 4 °C. Frozen liquid store at -20°C to -80°C. Recommended to avoid repeate e-and-thaw cycles.	Storage	Lyophilized powder store at 4 °C. Frozen liquid store at -20°C to -80°C. Recommended to avoid repeated freeze- e-and-thaw cycles.