



## Lyophilized DOTAP/DOPC/CHOL/DSPE-mPEG2000 ATP Liposome, NBD-lipid Labeled

Catalog: Lipo-246RG

## PRODUCT INFORMATION

Name	Lyophilized DOTAP/DOPC/CHOL/DSPE-mPEG2000 ATP Liposome, NBD-lipid Labeled
Cat No.	Lipo-246RG
Product Overview	The encapsulation of ATP in liposomes markedly promotes its effectiveness by preventing the hydrolysis by extracellular enzymes, increasing ATP circulation time and enhancing its intracellular penetration. ATP liposomes can be used in various models such as myocardial, liver, retina and wound healing ischemia. Studies have sown the ability of liposomal encapsulated ATP to prevent cell death and tissue dysfunction following ischemic events. The concentration of encapsulated ATP is 0.5µmol/vial. Creative Biostructure could customize different DOTAP/DOPC/CHOL ratio to meet your requirements. We can also manufacture empty lyophilized liposomes (without ATP) for control with the same lipid composition as your desired.
Lipid Composition	DOTAP/DOPC/CHOL/DSPE-mPEG2000/ NBD-PE (0.9/0.075/0.435/0.075/0.015 µmol/vial) DOTAP: 1,2-dioleoyl-3-trimethylammonium-propane (chloride salt) DOPC: 1,2-dioleoyl-sn-glycero-3-phosphocholine CHO L: Cholesterol NBD PE: 1,2-dioleoyl-sn-glycero-3-phosphoethanolamine-N-(7-nitro-2-1,3-benzoxadiazol-4-yl) (ammonium salt) (NBD PE) DSPE-mPEG2000: 1,2-distearoyl-sn-glycero-3-phosphoethanolamine-N-[methoxy(polyethyleneglycol)-2000] (ammonium salt)/CAS: 474922-77-5
Form	Lyophilized Powder
Storage Buffer	PBS, pH 7.4 with trehalose as lyoprotectant
Concentration	Lipid Concentration 1.5 µmol/vial
Stability	6 months
Storage	-20°C